South Australian Perinatal Practice Guideline

# Hypertensive Disorders in Pregnancy

© Department for Health and Wellbeing, Government of South Australia. All rights reserved.

Flowchart 3: Initial Management of Eclampsia[2]

if6-1342(eo)-1652(n P)2Span<</MCID 051

## Summary of Practice Recommendations

Women with an upper arm circumference > 33 cm require an alternate cuff
Women should be assessed for risk factors and commenced on prophylaxis if appropriate

## Definitions

| Term | Definition |
|---|---|
| **Hypertension** | Systolic BP ≥ 140 mm Hg and/or diastolic BP ≥ 90 mmHg (K5)<br>Confirmed by repeated readings over several hours |
| **Severe hypertension** | Systolic BP ≥160 mm Hg and/or diastolic BP ≥ 110 mmHg |
| **Preeclampsia** | A multi-system disorder characterised by hypertension and involvement of one or more other organ systems and/or the fetus<br>Proteinuria is the most commonly recognised additional feature after hypertension (not mandatory for clinical diagnosis) |
| **Gestational hypertension** | New onset of hypertension after 20 weeks gestation without any maternal or fetal features of preeclampsia<br>Return of BP to normal within 3 months postpartum |
| **Chronic hypertension** | Includes essential hypertension as well as hypertension secondary to a range of conditions<br>**Essential hypertension:** A systolic BP ≥ 140 mmHg and/or a diastolic BP ≥ 90 mmHg before pregnancy or before 20 weeks gestation without a known cause.<br>It may include women presenting early in pregnancy on antihypertensive medications where no secondary cause for hypertension has been determined.<br>**Secondary hypertension** may be due to: chronic kidney disease, renal artery stenosis, systemic disease with renal involvement, endocrine disorders, coarctation of the aorta, medications<br>**White coat hypertension** refers to raised BP in the presence of a clinical attendant but normal BP otherwise as assessed by ambulatory or home BP monitoring |
| **Preeclampsia superimposed on chronic hypertension** | Pre-existing hypertension is a strong risk factor for the development of preeclampsia<br>**Superimposed preeclampsia** is diagnosed when chronic hypertension develops one or more of the systemic features of preeclampsia after 20 weeks of gestation<br>Substantial increases in proteinuria and hypertension should raise suspicion of preeclampsia |

## Introduction

This PPG is based on the recommendations from a multidisciplinary working party convened by the Society of Obstetric Medicine of Australia and New Zealand (SOMANZ)[1] along with management principles outlined in the Practical Obstetric Multi-Professional Training (PROMPT) course manual[2] .

### Recording blood pressure in pregnancy

| Technique | Procedure | Rationale |
|---|---|---|
| **Position of woman** | Seated comfortably, legs resting on a flat surface and her arm resting at heart level | Different arm positions can produce significantly different measurements |
| | Measure BP on both arms at initial antenatal visit | |
| | In labour, measure in lateral recumbency | Avoid supine hypotension |
| | Avoid supine posture | |
| **Cuff size** | **Upper arm circumference 33-44 cm** | |
| | Use a large cuff with an inflatable bladder covering 80% of the arm circumference | |
| | **Upper arm circumference > 44 cm** | |

## Diagnosis of preeclampsia

A diagnosis of preeclampsia can be made when hypertension arises after 20 weeks gestation and is accompanied by one or more of the following signs of organ involvement:

| | |
|---|---|
| **Renal** | Significant proteinuria – either 300 mg per 24 hours or a spot urine protein/creatinine ratio of 30 mg/mmol or more<br>Serum or plasma creatinine 90 micromols/L or more<br>Oliguria: less than 80 mL / 4 hours<br>Urate is not included as a diagnostic feature |
| **Haematological** | Thrombocytopenia less than100,000 /microlitre<br>Haemolysis: schistocytes or red cell fragments on blood film, raised bilirubin, raised lactate dehydrogenase more than 600 international units/L, decreased haptoglobin<br>Disseminated intravascular coagulation |
| **Liver** | Raised serum transaminases<br>Severe epigastric and / or right upper quadrant pain |
| **Neurological** | Convulsions (automatically redefines it as eclampsia)<br>Hyperreflexia with sustained clonus<br>Persistent, new headache<br>Persistent visual disturbances (photopsia, scotomata, cortical blindness, |

## Treatment of severe preeclampsia

Severe preeclampsia has been defined as BP ≥160/110 mmHg with proteinuria (urinary protein:creatinine ratio > 30mg/mmol or 24 hour urinary protein > 300 mg) **OR**

BP 140/90 – 159/109 mmHg with proteinuria with at least one of the following:

- Severe headache
- Visual disturbances
- Severe pain just below the ribs or vomiting
- Papilloedema
- Signs of clonus (≥ 3 beats)
- Liver tenderness
- HELLP syndrome
- Platelet count < 100 x 109/L
- Abnormal liver enzymes

There are three main aspects to care of the woman with severe preeclampsia (see flowchart 2):

**1. Stabilise**

- Control BP
- Prevent seizures using Magnesium Sulphate IV (see appendix 3)

**2. Monitor**

- Vital signs
- Urinary output
- Neurological status
- Clotting factors
- Fetal condition

**3. Plan for labour/birth**

## Labetalol intravenous infusion

If the blood pressure is not adequately controlled after 2 bolus doses, consider a continuous labetalol infusion, either via a syringe driver or an infusion pump. Max. dose = 300 mg/24 hrs

| Syringe driver | Infusion pump |
|---|---|
| **Set up**<br>> Draw up labetalol 200 mg (40 mL) undiluted<br>> Using medication added sticker write "labetalol 5 mg in 1 mL" and attach label to syringe | **Set up**<br>> Withdraw 40 mL from a 100 mL bag of sodium chloride 0.9 %<br>> Draw up 200 mg labetalol (40 mL) and add to remaining 60 mL in the bag of sodium chloride 0.9 %<br>> Using medication added sticker write "labetalol 2 mg in 1 mL in sodium chloride 0.9 % (labetalol 200 mg made up to 100 mL with sodium chloride 0.9 %)" and attach label to bag |
| **Syringe driver infusion dose**<br>> Start infusion at 4 mL / hour (20 mg / hour). Titrate to stabilise blood pressure by maintaining or adjusting (e.g. doubling, halving) the infusion as required every 15-30 minutes to a maximum dose of 32 mL / hour (160 mg / hour)<br>> Discontinue by weaning over 1-2 hours when blood pressure is consistently less than155/95 mmHg | **Infusion pump dose**<br>> Start infusion at 10 mL / hour (20 mg / hour). Titrate to stabilise blood pressure by maintaining or adjusting (e.g. doubling, halving) the infusion as required every 15-30 minutes to a maximum dose of 80 mL / hour (160 mg / hour)<br>> Discontinue by weaning over 1-2 hours when blood pressure is consistently less than155/95 mmHg |

**Observations**

> Measure blood pressure and pulse every 15-30 minutes until stabilised, then record every hour as required
  - >If blood pressure decreases precipitously, halve the infusion rate or cease (depending on severity)
  - >Blood pressure should not be lowered below 140/85 mmHg
> Continuous electronic fetal monitoring during intravenous administration

### Relative contraindications

> Bronchial asthma or chronic obstructive airways disease
> For more contraindications refer to the Product Information or the Australian Medicines Handbook[3]

### Side effects

> Hypotension: cease if blood pressure < 140 mm Hg systolic
> Bradycardia: cease if heart rate < 60/minute
> Fetal bradycardia
> Wheezing and bronchospasm: cease if severe
> Headache and nausea
> Extravasation of labetalol solution may cause tissue damage. Stop and seek urgent assistance to re-site the IV
> Blurred vision and/or retention of urine may occasionally be seen, as may scalp tingling that may last up to 24-48 hours
> Prolonged maternal high doses of labetalol may cause hypotension in the preterm growth restricted newborn

| Hydralazine IV infusion administration | |
|---|---|
| **Administration precautions** | > **A precipitous fall in blood pressure can occur after intravenous hydralazine which may impair placental perfusion resulting in fetal distress.**<br>> If there is a risk of hypovolaemia, give intravenous fluid preload of 250 mL of either sodium chloride 0.9 % or Hartmann's immediately before use – see Fluid Management<br>> May be administered by a midwife under the supervision of a medical officer |
| **Intravenous Infusion (syringe pump)** | > **Administer via syringe pump**<br>> Hydralazine 1 mg/mL (1 in 1000) |

## Appendix 3: Magnesium Sulphate Infusion Regimen & Intramuscular Dose

## Magnesium sulphate volumetric infusion pump regimen

**Note: A volumetric infusion pump should only be utilised for the administration of magnesium sulphate where there is no access to a syringe driver**

### Administration precautions

> Magnesium sulphate is a High Risk Medicine

> Use a dedicated intravenous line for magnesium sulphate .

### Relative contraindications

**The use of this drug can be hazardous in association with:**

> Renal failure or severe renal compromise

> Hypocalcaemic states

> Other drugs, especially vasta 0 8TJ 0 Tc 0 Tw 12. ( s)696.4 (ugs)-8 (0 Tc 0 Tw 12.42 p3 (r)-1.dLBoMC /LB

## Acknowledgements

The South Australian Perinatal Practice Guidelines gratefully acknowledge the contribution of clinicians and other stakeholders who participated throughout the guideline development process particularly:

**Write Group Leads**

## Document Ownership & History

**Developed by:** SA Maternal, Neonatal & Gynaecology Community of Practice
**Contact:**