OFFICIAL

SAPPG Upload to SA Health Website and Practice Guidelines Web-Based App Summary

Date uploaded: 17/04/2024

Title of Document (previous title if applicable)	Revised or New	Summary of key points / changes	Risk management concerns +/- budget implications
Postpartum Haemorrhage	Revised (version 6)	This PPG has been reviewed in line with current international and national guidelines, and latest research recommendations for the management of PPH.	Distribution of information to all
Note that the following PPGs have been merged into the PPH PPG and removed from circulation: • Ergot derivatives • Prostaglandin analogues • Secondary PPH • Intra-uterine balloon tamponade • Oxytocin prophylaxis for third stage management and PPH		Key points: • Use of Carbetocin for active management of third stage of labour has been added. To be used in vaginal births only when women meet the criteria of: • PPH risk factors and/or p resence of con/cardiac anomalies. Note: Carbetocin is not currently on the SA formulary for vaginal births therefore an IPU form must be completed and sent to pharmacy. • Addition of tranexamic acid as part of first line PPH management medication • Addition of Carboprost as part of second line PPH management medication • New primary PPH management algorithm that includes tranexamic acid and carboprost. • External link for larger, printable flowchart – access via the link at the bottom of page 2 of the PPG. Print as needed. • Secondary PPH management flowchart updated with improved readability. • All blood loss must be measured (e.g., weighing soiled linen/pads). • Previous reoddsmetidalish tabless carbost at account of page 2 of the PPG with table updated. • PPH record form updated to reflect new PPH management flowchart. • Suggested citation added (New) • Appendices updated.	ion no lo)rrTw 6 Tf0 Tc -25⁄188g– ∮8∮